

# Basic Differentiation - A Refresher

v3. March 2003



www.mathcentre.ac.uk

 $\bigcirc$  mathcentre 2003

## Contents

	Foreword	2
	Preliminary work	2
	How to use this booklet	2
	Reminders	3
	Introduction	4
1.	Differentiation of a simple power	5
2.	Differentiation of a unit power	6
3.	Differentiation of a constant	7
4.	Differentiation of a simple power multiplied by a constant	8
	Notes page	9
5.	Differentiation of a unit power multiplied by a constant	10
6.	Differentiation of a general power	11
7.	Differentiation of a general power multiplied by a constant	12
8.	Differentiation of a sum or difference of terms	13
	Notes page	14
9.	Differentiation of a simple fraction	15
10.	Differentiation of fractions reducible to simpler fractions	16
	Notes page	17
11.	Differentiation of a simple bracket	18
12.	Differentiation of a general bracket	19
	Notes page	20
13.	Miscellaneous problems	21
	Answers to problems	24
	Acknowledgements	28

## Foreword

The material in this refresher course has been designed to enable you to cope better with your university mathematics programme. When your programme starts you will find that the ability to differentiate confidently will be invaluable. We think that this is so important that we are making this course available for you to work through either before you come to university, or during the early stages of your programme.

## **Preliminary work**

You are advised to work through the companion booklet An Algebra Refresher before embarking upon this basic differentiation revision course.

### How to use this booklet

You are advised to work through each section in this booklet in order. You may need to revise some topics by looking at an AS-level textbook which contains information about differentiation.

You should attempt a range of questions from each section, and check your answers with those at the back of the booklet. The more questions that you attempt, the more familiar you will become with these vital topics. We have left sufficient space in the booklet so that you can do any necessary working within it. So, treat this as a work-book.

If you get questions wrong you should revise the material and try again until you are getting the majority of questions correct.

If you cannot sort out your difficulties, do not worry about this. Your university will make provision to help you with your problems. This may take the form of special revision lectures, self-study revision material or a drop-in mathematics support centre.



2

## Reminders

Use this page to note topics and questions which you found difficult.

Seek help with these from your tutor or from other university support services as soon as possible.



## Introduction

Differentiation is an aspect of calculus that enables us to determine how one quantity changes with regard to another.

In this booklet we will not however be concerned with the applications of differentiation, or the theory behind its development, but solely with the mechanisms by which the process of differentiation may be quickly and effectively carried out.

Each section that follows will introduce a particular technique followed by several problems for you to practice what you have learned.

At the end of the booklet you will find answers for each of the sections.

Included are some pages for you to make notes that may serve as a reminder to you of any possible areas of difficulty. You should seek help with such areas of difficulty from your tutor or other university support services.



## 1. Differentiation of a simple power

To differentiate $s = t^n$ .	Example
• Bring the existing power down and use it to multiply.	$s = t^3$
• Reduce the old power by one and use this as the	$\frac{ds}{dt} = 3t^{3-1}$
new power. <b>Answer</b> $\frac{ds}{dt} = nt^{n-1}$	$= 3t^2$

**Practice:** In the space provided write down the requested derivative for each of the following expressions.

(a) $s = t^9$ ,	$\frac{ds}{dt}$ (b)	$v = t^4$ ,	$\frac{dv}{dt}$ (c)	$a = t^5$ ,	$\frac{da}{dt}$	(d) $y = x^7$ ,	$\frac{dy}{dx}$
(e) $y = x^{12}$ ,	$\frac{dy}{dx}$ (f)	$V = r^3$ ,	$\frac{dV}{dr}$ (g)	$P = v^6$ ,	$\frac{dP}{dv}$	(h) $A = r^2$ ,	$\frac{dA}{dr}$
(i) $T = \theta^8$ ,	$\frac{dT}{d\theta}$ (j)	$R = t^{19},$	$\frac{dR}{dt}$ (k)	$v = t^{11},$	$\frac{dv}{dt}$	(l) $p = V^{10}$ ,	$\frac{dp}{dV}$

5

## 2. Differentiation of a unit power

To differentiate s = t.

- The phrase 'a unit power' refers to the fact that the power is 1.
- Note that t is the same as  $t^1$ . We can use the previous rule.
- Differentiating the term  $t = t^1$  gives  $1t^0 = 1$

**Answer**  $\frac{ds}{dt} = 1$ 

Practice:	In the space provided write down the requested derivative for each of
	the following expressions.

(a) $r = t$ ,	$\frac{dr}{dt}$ (b) $v = t$ ,	$\frac{dv}{dt}  (c)  a = t,$	$\frac{da}{dt}  (d)  y = x,$	$\frac{dy}{dx}$
(e) $z = y$ ,	$\frac{dz}{dy}  (f)  V = r,$	$\frac{dV}{dr}  (g)  P = v,$	$\frac{dP}{dv}  (h)  W = z,$	$\frac{dW}{dz}$
(i) $T = \theta$ ,	$\frac{dT}{d\theta}  (j)  Q = r,$	$\frac{dQ}{dr}  (\mathbf{k})  m = s,$	$\frac{dm}{ds}  (1)  x = y,$	$\frac{dx}{dy}$

## 3. Differentiation of a constant

To differentiate $s = a$ where $a$ is a constant.	Example
• An isolated constant always differentiates to zero.	s = 4
<b>Answer</b> $\frac{ds}{dt} = 0$	$\frac{ds}{dt} = 0$

**Practice:** In the space provided write down the requested derivative for each of the following expressions.

(a) $s = 5$ ,	$\frac{ds}{dt}  \text{(b)}  v = 0.6,$	$\frac{dv}{dt}  (c)  \theta = \frac{1}{2},$	$\frac{d\theta}{dt}$ (d) $r = 1$ , $\frac{dr}{d\theta}$	
(e) $T = \pi$ ,	$\frac{dT}{dt}  \text{(f)}  z = 29.6,$	$\frac{dz}{dx}  (g)  x = 3^{-1},$	$\frac{dx}{dt}$ (h) $f = 2.7,  \frac{df}{dx}$	
(i) $p = 4$ ,	$\frac{dp}{dr}$ (j) $q = \sqrt{2}$ ,	$\frac{dq}{dt}  (k)  w = \frac{3}{4},$	$\frac{dw}{dz}$ (l) $m = 2.7,  \frac{dm}{d\theta}$	

## 4. Differentiation of a simple power multiplied by a constant

To differentiate $s = at^n$ where a is a constant.	Example
• Bring the existing power down and use it to multiply.	Example $s = 3t^4$
• Reduce the old power by one and use this as the new power. ds $ds$ $ds$ $ds$ $ds$ $ds$ $ds$ $ds$	$\frac{ds}{dt} = 4 \times 3t^{4-1}$ $= 12t^3$
<b>Answer</b> $\frac{ds}{dt} = ant^{n-1}$	$= 12t^{3}$

(a) $s = 3t^4$ ,	$\frac{ds}{dt}$	(b) $y = 7x^3$ ,	$\frac{dy}{dx}$	(c) $r = 0.4\theta^5$ ,	$\frac{dr}{d\theta}$	(d) $v = 4r^3$ ,	$\frac{dv}{dr}$
(e) $A = \pi r^2$ ,	$\frac{dA}{dr}$	(f) $T = 6\theta^4$ ,	$\frac{dT}{d\theta}$	(g) $v = 5t^9$ ,	$\frac{dv}{dt}$	(h) $p = 6v^3$ ,	$\frac{dp}{dv}$
(i) $y = 14x^2$ ,	$\frac{dy}{dx}$	(j) $r = 5t^7$ ,	$\frac{dr}{dt}$	(k) $P = 7v^6$ ,	$\frac{dP}{dv}$	(l) $C = 12N^8$ ,	$\frac{dC}{dN}$

## Notes Page



9

## 5. Differentiation of a unit power multiplied by a constant

To differentiate $s = at$ where $a$ is a constant.	Example
• The result is always the same as the constant.	s = 3t
<b>Answer</b> $\frac{ds}{dt} = a$	$\frac{ds}{dt} = 3$

Practice: In the space provided write down the requested derivative for each of the following expressions.

(a) $y = 5x$ ,	$\frac{dy}{dx}$	(b) $s = 6t$ ,	$\frac{ds}{dt}$ (c)	p = 4v,	$rac{dp}{dv}$
(d) $C = 7n$ ,	$\frac{dC}{dn}$	(e) $Q = 0.5v$ ,	$\frac{dQ}{dv} \qquad (f)$	$z = \pi w,$	$\frac{dz}{dw}$
		(h) $\theta = \frac{1}{3}t,$	$\frac{d\theta}{dt}$ (i)	$r = \frac{1}{4}t,$	$\frac{dr}{dt}$
(j) $s = 3.14\omega$ ,	$\frac{ds}{d\omega}$	(k) $w = 5z$ ,	$\frac{dw}{dz} \qquad (1)$	$R = 9.6\phi,$	$\frac{dR}{d\phi}$



## 6. Differentiation of a general power

To differentiate $s = t^n$ for any value of $n$ .	Example
• Bring the existing power down and use it to multiply.	$s = t^{-3.4}$
• Reduce the old power by one and use this as the new power.	$\frac{ds}{dt} = -3.4t^{-3.4-1}$
<b>Answer</b> $\frac{ds}{dt} = nt^{n-1}$	$= -3.4t^{-4.4}$

(a) $s = t^{2.8}$ ,	$\frac{ds}{dt}$	(b) $y = x^{2/3}$ ,	$\frac{dy}{dx}$	(c) $r = 0.4\theta^{1/4}$ ,	$\frac{dr}{d\theta}$
(d) $v = 4r^{-3}$ ,	$\frac{dv}{dr}$	(e) $A = r^{1/2}$ ,	$\frac{dA}{dr}$	(f) $T = \theta^{-6}$ ,	$\frac{dT}{d\theta}$
(g) $v = t^{-1}$ ,	$\frac{dv}{dt}$	(h) $p = v^{-3.5}$ ,	$\frac{dp}{dv}$	(i) $y = x^{3/4}$ ,	$\frac{dy}{dx}$
(j) $r = t^{-1/3}$ ,	$\frac{dr}{dt}$	(k) $P = v^{1.8}$ ,	$\frac{dP}{dv}$	(l) $C = N^{-6}$ ,	$\frac{dC}{dN}$

## 7. Differentiation of a general power multiplied by a constant

To differentiate $s = at^n$ for any value of $n$ where $a$ is a constant.	
• Bring the existing power down and use it to multiply.	$s = 2.5t^{-4}$
• Reduce the old power by one and use this as the new power.	$\frac{ds}{dt} = -4 \times 2.5t^{-4-1}$
<b>Answer</b> $\frac{ds}{dt} = ant^{n-1}$	$= -10t^{-5}$

(a) $y = 5x^{1.5}$ ,	$\frac{dy}{dx}$	(b) $s =$	$4t^{-3},$	$\frac{ds}{dt}$	(c)	$p = 8v^{1/2},$	$\frac{dp}{dv}$
(d) $C = 6n^{-1}$ ,	$\frac{dC}{dn}$	(e) $Q =$	$= 2v^{0.5},$	$\frac{dQ}{dv}$	(f)	$z = 6w^{-2.5},$	$\frac{dz}{dw}$
(g) $N = 5.2n^{10}$ ,	$\frac{dN}{dn}$	(h) $\theta =$	$=\frac{1}{5}t^{6},$	$\frac{d\theta}{dt}$	(i)	$r = -5t^{-2},$	$\frac{dr}{dt}$
(j) $s = 2.5p^{2.5}$ ,	$\frac{ds}{dp}$	(k) <i>w</i> =	$= 15z^{-0.1},$	$\frac{dw}{dz}$	(l)	$v = -3r^{-4},$	$\frac{dv}{dr}$

### 8. Differentiation of a sum or difference of terms

T

To different	tiate $s = f(t) + g(t)$ or $s = f(t) - g(t)$ .	Example		
• Differenti	ate $f(t)$ and $g(t)$ using previous rules.	$s = 3t^2 \pm 6t^{\frac{1}{3}}$		
• Add or su	ibtract the results.	$\frac{ds}{dt} = 2 \times 3t^{2-1} \pm \frac{1}{3} \times 6t^{\frac{1}{3}-1}$		
Answer	$\frac{ds}{dt} = \frac{df}{dt} + \frac{dg}{dt}$ or $\frac{ds}{dt} = \frac{df}{dt} - \frac{dg}{dt}$	$= 6t \pm 2t^{-\frac{2}{3}}$		

(a) $s = t^2 + t^3$ ,	$\frac{ds}{dt}$	(b) <i>v</i>	$v = 2t + 3t^4,$	$\frac{dv}{dt}$	(c)	a = 3 + 4t,	$\frac{da}{dt}$
(d) $v = 5t - 10t$	$\frac{dv}{dt}$	(e) <i>a</i>	$= 3t^{4.5} - 4t^{-1}$	$\frac{da}{dt}$	(f)	$s = 4t^{1/2} - 2.5t^{-2},$	$\frac{ds}{dt}$
(g) $y = x^2 + 3x^2$	$\frac{dy}{dx}$	(h) <i>y</i>	$y = 2x^{-1} - x^{-2},$	$\frac{dy}{dx}$	(i)	$x = 3y + \frac{1}{4}y^2,$	$\frac{dx}{dy}$



## Notes Page



## 9. Differentiation of a simple fraction

To differentiate $s = \frac{a}{t^n}$ where a is a constant.	Example
• Rewrite as $s = at^{-n}$ .	$s = \frac{2}{t^3}$
	$= 2t^{-3}$
• Differentiate using previous rules.	$s = \frac{2}{t^3}$ $= 2t^{-3}$ $\frac{ds}{dt} = -3 \times 2t^{-3-1}$
<b>Answer</b> $\frac{ds}{dt} = -ant^{-n-1}$	$= -6t^{-4}$

Practice:	In the space provided write down the requested derivative for each of
	the following expressions.

(a) $s = \frac{1}{t}$ ,	$\frac{ds}{dt}$	(b) $s = \frac{2}{t^2},$	$\frac{ds}{dt}$	(c) $y = \frac{5}{x^3}$ ,	$\frac{dy}{dx}$
(d) $p = \frac{2.4}{r^7},$	$\frac{dp}{dr}$	(e) $v = \frac{4.5}{t^{1/2}},$	$\frac{dv}{dt}$	(f) $a = \frac{10}{t^6}$ ,	$\frac{da}{dt}$
(g) $v = \frac{4.9}{x^4},$	$\frac{dv}{dx}$	(h) $s = \frac{2.5}{\theta^3},$	$\frac{ds}{d\theta}$	(i) $x = \frac{\pi}{\theta}$ ,	$\frac{dx}{d\theta}$
(j) $\theta = \frac{\sqrt{2}}{t^5},$	$\frac{d\theta}{dt}$	$(\mathbf{k})  z = \frac{7}{2x},$	$\frac{dz}{dx}$	(l) $q = \frac{3}{r^{2.5}},$	$\frac{dq}{dr}$

# 10. Differentiation of fractions reducible to simpler fractions

To differentiate $s = \frac{a + bt^n}{ct^m}$ .	Example
• Rewrite as $s = \frac{a}{ct^m} + \frac{bt^n}{ct^m}$ .	$s = \frac{4 + 3t^2}{5t^3}$ $= \frac{4}{5t^3} + \frac{3t^2}{5t^3}$
• Use power laws to simplify.	$=\frac{4}{5t^3} + \frac{3t^2}{5t^3}$
• Differentiate using previous rules.	$=\frac{4t^{-3}}{5}+\frac{3t^{-1}}{5}$
	$\frac{ds}{dt} = -\frac{3 \times 4t^{-3-1}}{5} + (-1) \times \frac{3t^{-1-1}}{5}$
Answer $\frac{ds}{dt} = -\frac{amt^{-m-1}}{c} + \frac{b}{c}(n-m)t^{n-m-1}$	$= -\frac{12t^{-4}}{5} - \frac{3t^{-2}}{5}$

**Practice:** In the space provided write down the requested derivative for each of the following expressions.

(a) $s = \frac{4t+1}{t}$ ,	$\frac{ds}{dt}$	(b)	$s = \frac{5t^3 + 3t}{t},$	$\frac{ds}{dt}$	(c)	$v = \frac{5t^2 + 3t + 1}{t^3},$	$\frac{dv}{dt}$
(d) $y = \frac{7+6x+5x^2}{x^2}$ ,	$\frac{dy}{dx}$	(e)	$y = \frac{3+5x}{4x},$	$\frac{dy}{dx}$	(f)	$z = \frac{7 + 2t}{3t},$	$\frac{dz}{dt}$
(g) $\theta = \frac{2+5t+2t^3}{t^2},$	$\frac{d\theta}{dt}$	(h)	$w = \frac{8+6z^2}{3z^4},$	$\frac{dw}{dz}$	(i)	$x = \frac{3y + 2y^2}{4y},$	$\frac{dx}{dy}$



## Notes Page



17

# 11. Differentiation of a simple bracket

To differentiate $s = (at + b)^n$ .	Example
• Bring the existing power down and use it to multiply.	$s = (3t+6)^5$
• Reduce the old power by 1 and use this as the new power.	
• Differentiate the expression inside the bracket.	
• Multiply your results together.	$\frac{ds}{dt} = 5 \times (3t+6)^{5-1} \times 3$
<b>Answer</b> $\frac{ds}{dt} = an(at+b)^{n-1}$	$= 15(3t+6)^4$

(a) $s = (2t+3)^6$ ,	$\frac{ds}{dt}$	(b)	$v = (5+2t)^7,$	$\frac{dv}{dt}$	(c)	$\theta = (0.4t + 1)^{2.5},$	$\frac{d\theta}{dt}$
(d) $z = (\pi x + 2.4)^{10}$ ,	$\frac{dz}{dx}$	(e)	$a = (5t - 3)^{-2},$	$\frac{da}{dt}$	(f)	$s = (3 - 6t)^{-3},$	$\frac{ds}{dt}$
(g) $y = (4x - 1)^{1/2}$ ,	$\frac{dy}{dx}$	(h)	$y = (2 + 0.5t)^3,$	$\frac{dy}{dt}$	(i)	$z = (4x - 6)^{-0.5},$	$\frac{dz}{dx}$

## 12. Differentiation of a general bracket

To differentiate $s = [f(t)]^n$ .	Example
• Bring the existing power down	$s = (5t^3 - 3t^2 + 1)^7$
and use it to multiply.	
$\bullet$ Reduce the old power by 1	
and use this as the new power.	
• Differentiate $f(t)$ to give $\frac{df}{dt}$ .	
• Multiply your results together.	$\frac{ds}{dt} = 7 \times (5t^3 - 3t^2 + 1)^{7-1} \times (3 \times 5t^{3-1} - 2 \times 3t^{2-1} + 0)$
• Simplify.	$= 7(5t^3 - 3t^2 + 1)^6(15t^2 - 6t)$
<b>Answer</b> $\frac{ds}{dt} = n[f(t)]^{n-1} \cdot \frac{df}{dt}$	$= 21t(5t^3 - 3t^2 + 1)^6(5t - 2)$

# **Practice:** In the space provided write down the requested derivative for each of the following expressions.

(a) $s = (3t^2 + 2t + 4)^5$	$\frac{ds}{dt}$	(b) $r = (4\theta^2 - 2\theta + 1)^{-2}$	$\frac{dr}{d\theta}$	(c) $v = (8x^2 + 5x)^2$	$\frac{dv}{dx}$
(d) $\theta = (3t+1)^{-1}$		(e) $p = (5v^2 + 2)^{-2}$		(f) $y = (t^3 + t + 1)^2$	
(g) $z = (5x^3 + 1)^{-4}$	$\frac{dz}{dx}$	(h) $s = (2t+1)^4$	$\frac{ds}{dt}$	(i) $x = (3y^2 - 4y)^3$	$\frac{dx}{dy}$



Notes Page



## 13. Miscellaneous problems


1. In each of the following determine 
$$\frac{ds}{dt}$$
.

$$\begin{array}{lll} (a) & s = \frac{2}{t} & (b) & s = t^{1/2} & (c) & s = 4t^{-2} \\ (d) & s = \frac{1}{3}t & (e) & s = \frac{2t+1}{t} & (f) & s = \frac{t^3-3t+4}{t^2} \\ (g) & s = (5t+1)^4 & (h) & s = (3t+1)^{-1} & (i) & s = \frac{2}{3t+1} \\ (j) & s = 5t^2+3t+6 & (k) & s = 2t^{-3}+t^4 & (l) & s = (2t+3t^{-1})^{-5} \end{array}$$



2. In each of the following determine  $\frac{dy}{dx}$ .

(a) $y = x^{1/2}$	(b) $y = (x+1)^{1/2}$	$(c)  y = \frac{3}{x} + 1$
$(d)  y = 5x + \frac{1}{x}$	$(e)  y = 5x^2$	$(f)  y = 7(x^2 + 1)^5$
(g) $y = (2x+1)^{1/3}$	$(h)  y = \frac{7+2x}{x}$	$(i)  y = \frac{5}{3x^2}$
$(j)  y = 4x^{-2} + 3x^{-1}$	$(k)  y = 24 - 2x^{-3/4}$	(l) $y = (2x^3 + 3x^2 - 5x)^2$





3. In each of the following determine  $\frac{dv}{dt}$ .

(a) $v = 3t$	(b) $v = 0.7t^{-3} - 6t^3$	(c) $v = (5t - 6)^{-0.5}$
$(d)  v = \frac{4}{t} + 3t$	(e)  v = 6	$(f)  v = 2t^{-2} - 3t^{-1} + 4 + 5t$
$(g)  v = (6t - 2)^{-0.5}$	$(h)  v = \frac{3}{t^2}$	( <i>i</i> ) $v = 3t - \frac{1}{t^{0.1}}$
$(j)  v = 8t^4 - 3t^{1/2}$	$(k)  v = (5t^{-2} + 3t^2)^3$	$(l)  v = \pi - \pi t$



(d)  $7x^6$ 

 $(h) \quad 2r$ 

 $(l) \quad 10V^9$ 

## Answers to Problems

(c)  $5t^4$ 

 $(g) \quad 6v^5$ 

 $(k) \quad 11t^{10}$ 

# Section 1 (a) $9t^8$ (e) $12x^{11}$ (i) $8\theta^7$

### Section 2

(a) 1	(b) 1	(c) 1	(d) 1
(e) 1	(f) 1	(g) 1	(h) 1
(i) 1	(j) 1	(k) 1	(l) 1

(b)  $4t^3$ 

 $(f) \ 3r^2$ 

(j) 19 $t^{18}$ 

### Section 3

(a)	0	(b)  0	(c) 0	(d) 0
(e)	0	(f)  0	(g)  0	(h)  0
(i)	0	(j) 0	(k) 0	(l)  0

### Section 4

(a)	$12t^{3}$	(b) $21x^2$	$(c)$ $2\theta^4$	(d)	$12r^2$
(e)	$2\pi r$	$(f)  24\theta^3$	$(g)  45t^8$	(h)	$18v^2$
(i)	28x	$(j)$ 35 $t^6$	$(k)$ 42 $v^5$	(l)	$96N^7$

### Section 5

(a) 5	(b) 6	(c) 4	(d) 7
(e)  0.5	$(f)$ $\pi$	(g) 1.7	( <i>h</i> ) $1/3$
( <i>i</i> ) $1/4$	(j) 3.14	(k) 5	(l) 9.6



### Section 6

(a) $2.8t^{1.8}$	$(b)  \frac{2}{3}x^{-1/3}$	(c) $0.1\theta^{-3/4}$	$(d) - 12r^{-4}$
$(e)  \frac{1}{2}r^{-1/2}$	$(f) - 6\theta^{-7}$	$(g) - t^{-2}$	$(h) - 3.5v^{-4.5}$
$(i)  \frac{3}{4}x^{-1/4}$	$(j) - \frac{1}{3}t^{-4/3}$	$(k)$ 1.8 $v^{0.8}$	$(l) - 6N^{-7}$

### Section 7

(a)	$7.5x^{0.5}$	(b)	$-12t^{-4}$	(c)	$4v^{-1/2}$	(d)	$-6n^{-2}$
(e)	$v^{-0.5}$	(f)	$-15w^{-3.5}$	(g)	$52n^{9}$	(h)	$\frac{6}{5}t^{5}$
(i)	$10t^{-3}$	(j)	$6.25p^{1.5}$	(k)	$-1.5z^{-1.1}$	(l)	$12r^{-5}$

### Section 8

(a)	$2t + 3t^2$	(b) $2 + 12t^3$	(c) 4	(d) $5 - 20t$
(e)	$13.5t^{3.5} + 4t^{-2}$	$(f)  2t^{-1/2} + 5t^{-3}$	$(g)  2x + 12x^3$	$(h)  -2x^{-2} + 2x^{-3}$
(i)	$3 + \frac{y}{2}$			

### Section 9

(a)	$-t^{-2}$	$(b) - 4t^{-3}$	$(c) - 15x^{-4}$	(d)	$-16.8r^{-8}$
(e)	$-2.25t^{-3/2}$	$(f) - 60t^{-7}$	$(g) - 19.6x^{-5}$	(h)	$-7.5\theta^{-4}$
(i)	$-\pi\theta^{-2}$	$(j) - 5\sqrt{2}t^{-6}$	$(k)  -\frac{7}{2}x^{-2}$	(l)	$-7.5r^{-3.5}$

### Section 10

(a) $-t^{-2}$	(b) 10t	$(c)  -5t^{-2} - 6t^{-3} - 3t^{-4}$	$(d)  -14x^{-3} - 6x^{-2}$
(e) $-\frac{3}{4}x^{-2}$	$(f) - \frac{7}{3}t^{-2}$	$(g)  -4t^{-3} - 5t^{-2} + 2$	$(h)  -\frac{32}{3}z^{-5} - 4z^{-3}$
$(i)$ $\frac{1}{2}$			

www.mathcentre.ac.uk

© mathcentre 2003

### Section 11

(a)  $12(2t+3)^5$  (b)  $14(5+2t)^6$  (c)  $(0.4t+1)^{1.5}$  (d)  $10\pi(\pi x+2.4)^9$ (e)  $-10(5t-3)^{-3}$  (f)  $18(3-6t)^{-4}$  (g)  $2(4x-1)^{-1/2}$  (h)  $1.5(2+0.5t)^{2}$ (*i*)  $-2(4x-6)^{-1.5}$ 

### Section 12

### Section 13

#### Exercise 1

(a)  $-2t^{-2}$  (b)  $\frac{1}{2}t^{-1/2}$  $(c) - 8t^{-3}$ (e)  $-t^{-2}$  (f)  $1+3t^{-2}-8t^{-3}$ (d) 1/3 $(g) \quad 20(5t+1)^3 \qquad (h) \quad -3(3t+1)^{-2} \qquad (i) \quad -6(3t+1)^{-2}$ (j) 10t + 3 (k)  $-6t^{-4} + 4t^3$  (l)  $-5(2t + 3t^{-1})^{-6}(2 - 3t^{-2})$ 

#### Exercise 2

- (a)  $\frac{1}{2}x^{-1/2}$  (b)  $\frac{1}{2}(x+1)^{-1/2}$
- (e) 10x (f)  $70x(x^2+1)^4$ (d)  $5 - x^{-2}$
- $(g) \quad \frac{2}{3}(2x+1)^{-2/3} \qquad (h) \quad -7x^{-2}$
- $(j) 8x^{-3} 3x^{-2}$   $(k) \frac{3}{2}x^{-7/4}$

$$(c) - 3x^{-2}$$

$$(i) - \frac{10}{3}x^{-3}$$

(l) 
$$2(2x^3 + 3x^2 - 5x)(6x^2 + 6x - 5)$$

### Exercise 3

(a) 3(b) 
$$-2.1t^{-4} - 18t^2$$
(c)  $-2.5(5t - 6)^{-1.5}$ (d)  $-4t^{-2} + 3$ (e) 0(f)  $-4t^{-3} + 3t^{-2} + 5$ (g)  $-3(6t - 2)^{-1.5}$ (h)  $-6t^{-3}$ (i)  $3 + 0.1t^{-1.1}$ 

(j)  $32t^3 - 1.5t^{-1/2}$  (k)  $6(5t^{-2} + 3t^2)^2(-5t^{-3} + 3t)$  (l)  $-\pi$ 



### Acknowledgements

The materials in **Basic Differentiation - A Refresher** were prepared by David Pidcock at the Mathematics Education Centre, Loughborough University, on behalf of **math**centre.

The document was prepared in Latex by Joan de Souza for **math**centre.

Users are invited to send any suggestions for improvement to enquiries@mathcentre.ac.uk.

This edition of **Basic Differentiation - A Refresher** is published by **math**centre in March 2003.

If departments wish to customise and print their own copies, the contents of the book can be supplied in electronic format suitable for printing.

Further details from:

mathcentre c/o Mathematics Education Centre Loughborough University LE11 3TU

email: enquiries@mathcentre.ac.uk tel: 01509 227465 www: http://www.mathcentre.ac.uk

